

Pre-Procedure Check List for Dr. Edward Pang's Patients:

READ COMPLETELY, IF YOU DO NOT UNDERSTAND PLEASE ASK BEFORE YOUR PROCEDURE, FAILURE TO COMPLY MAY RESULT IN CANCELLATION OF YOUR PROCEDURE BECAUSE OF SAFETY ISSUES

Please **DO**:




- Take all blood pressure pills and heart medications prior to the procedure; your procedure will be **cancelled** if your blood pressure or other vital signs are unstable
- **Must** have a **driver** to take you home
- **Arrive** one (1) hour before the scheduled time, if you are not checked in at the Ambulatory Surgery Center by the above time, your procedure may be **cancelled**
- Tell us if you have **ANY allergies** to medications, including contrast dye, iodine or iodine containing substances, latex, or steroids. You will require special preparation prior to the procedure & if not disclosed until the day of the procedure we may need to reschedule
- If female, make sure that you are **not pregnant**; if there is ANY chance that you are pregnant or you are unsure – let us know **BEFORE** the procedure date. The X-Rays used during the procedure may harm your baby.
- Resume all **heart and blood pressure** medications after the procedure

Please **DO NOT**:



- You will require clearance from prescribing physician prior to stopping any blood thinning medications. See below for specific guidelines:
 - Aspirin or aspirin containing products: in general you may continue aspirin medication unless told otherwise. If needed, stop **6 days prior** to procedure and then restart **24 hours after** procedure.
 - NSAIDs (Ibuprofen, Naproxen, Meloxicam, Diclofenac, Motrin, Aleve, Celebrex etc): in general you may continue NSAID medications unless told otherwise. If needed, stop **1-4 days prior** to procedure and then restart **24 hours after** procedure.
 - Coumadin (Warfarin): stop **5 days prior** to procedure then may bridge with lovenox (see below guidelines for lovenox). Also will require INR to be < 1.2 prior to procedure. May restart **24 hours after** procedure.
 - LMWH (Lovenox): If for therapeutic treatment, stop **24 hrs prior** to procedure and restart **12-24 hrs after** procedure. If for prophylaxis, stop **12 hours prior** to procedure and restart **12-24hrs after** procedure.
 - Clopidogrel (Plavix): stop **7 days prior** to procedure, and restart **12- 24 hours after** procedure
 - Dabigatran (Pradaxa), Rivaroxaban (Xarelto), Apixaban (Eliquis): stop **3 – 4 (Pradaxa) days prior** to procedure and then restart **24 hours later**. Consider bridging with lovenox
- Fish Oil, Vitamin E, Ginko biloba: stop **5 days prior** to the procedure date
- **Do Not eat or drink 6 hours before the procedure**. If you require heart or blood pressure medications you may take these with sips of water at least 1 hours before the above arrival time



Specialty Surgical Center of Beverly Hills

8670 Wilshire Blvd Suite 300

Beverly Hills, CA 90211

Cross street: Willaman Dr.


Between Robertson Blvd and La Cienega Blvd

(310) 275-1646

www.sscbeverlyhills.com

Date of Procedure:

Arrival Time:



Precision Ambulatory Surgery Center

450 N. Roxbury Dr., Suite 240


Beverly Hills, CA 90210

Cross street: S. Santa Monica Blvd.

Between Santa Monica Blvd and Wilshire Blvd

(310) 651-2050

<https://beverlyhillsasc.com/pasc/>



Date of Procedure:

Arrival Time:



Advanced Surgical Center of Beverly Hills


8750 Wilshire Blvd. Suite 150

Beverly Hills, CA 90211

Cross street: N. Robertson Blvd

(424) 382-1137

<http://www.advancedsurgicalcenterofbh.com/>



Date of Procedure:

Arrival Time: